

2.2. What is the definition of the fluid?

- A) a substance that doesn't permanently resist distortion
- B) has shear stress
- C) a substance that does permanently resist distortion
- D) shear stress magnitudes depend on the volume of the fluid

3.1. What is the symbol for density?

- A)  $\beta$
- B)  $\rho$
- C)  $\eta$
- D)  $\sigma$

3.4. What is the unit of the relative density?

- A)  $\text{g/cm}^3$
- B)  $\text{N/m}^3$
- C) dimensionless
- D)  $\text{kg/m}^3$

3.7. Convert the density from kilogram/liter [ $\text{kg/L}$ ] to gram/cubic centimeter [ $\text{g/cm}^3$ ]!

11.22  $\text{kg/L} = \dots\dots\dots \text{g/cm}^3$

- A) 0.01122
- B) 11.22
- C) 11220
- D) 11220000

3.8. Convert the density from gram/cubic centimeter [ $\text{g/cm}^3$ ] to kilogram/cubic meter [ $\text{kg/m}^3$ ]!

2.9  $\text{g/cm}^3 = \dots\dots\dots \text{kg/m}^3$

- A) 2.9
- B) 0.0029
- C) 2900
- D) 2900000

3.12. Density of water is maximum at

- A)  $-4\text{ }^\circ\text{C}$
- B)  $4\text{ }^\circ\text{C}$
- C)  $100\text{ }^\circ\text{C}$
- D)  $0\text{ }^\circ\text{C}$

3.13. The calculation of the specific weight

- A)  $\gamma = \rho/g$
- B)  $\gamma = \rho\eta$
- C)  $\gamma = \rho v$
- D)  $\gamma = \rho g$

3.14. True or falls?

- A) Specific gravity usually means relative density with respect to water.
- B) The symbol of the relative density is R
- C) Relative density is dimensionless
- D) Relative density =  $\rho_{\text{substance}} / \rho_{\text{reference}}$
- E) Relative density is the specific weight


3.88. If the changes in density occurs at constant temperature, then the process is called

- A) adiabatic
- B) isothermal
- C) isobaric
- D) isochoric

3.89. In isochoric process .....is constant during process

- A) pressure
- B) temperature
- C) volume
- D) heat transfer

3.90. Consider the cycle in the diagram made of 2 isobaric process. What are these?



- A) AB and DC
- B) AD and DC
- C) BC and CD
- D) AD and BC

3.99. Isochoric process

- A) no change in temperature
- B) no change in volume
- C) no change in pressure
- D) the system does no work

3.103. Pressure law

- A)  $V / T = \text{constant}$
- B)  $p \cdot V = \text{constant}$
- C)  $p / T = \text{constant}$
- D)  $(p \cdot V) / T = n \cdot R$

3.104. Charles' law

- A)  $V / T = \text{constant}$
- B)  $p \cdot V = \text{constant}$
- C)  $p / T = \text{constant}$
- D)  $(p \cdot V) / T = n \cdot R$

3.105. The ideal gas equation

- A)  $V / T = \text{constant}$
- B)  $(p \cdot V) / T = n \cdot R$
- C)  $p \cdot V = \text{constant}$
- D)  $p / T = \text{constant}$

3.106. Boyle's law

- A)  $p / T = \text{constant}$
- B)  $V / T = \text{constant}$
- C)  $(p \cdot V) / T = n \cdot R$
- D)  $p \cdot V = \text{constant}$

3.107. True or false? Ideal gases.....

- A) collisions aren't elastic
- B) There are no intermolecular forces
- C) Molecules are in rapid motion
- D) The total volume of the molecules isn't negligible compared to the volume of the container

3.110. One kilogram of hydrogen is confined in a volume of 220 dm<sup>3</sup> at -55 °C. What is the pressure if R is 4.115 kJ/kg·K?

*Solution:*

$$p = \rho \times R \times T$$

$$\rho = m/V = 1/0.220 = 4.54 \text{ kg /m}^3$$

$$R = 4.115 \text{ kJ/kg}\cdot\text{K} = 4115 \text{ J/kg}\cdot\text{K}$$

$$T = -55 + 273 = 218 \text{ K}$$

$$p = 4.54 \times 4115 \times 218 = 4.073 \times 10^6 \text{ Pa}$$

3.111. In all gas equations, temperature is measured in

- A) Celsius
- B) Fahrenheit
- C) Kelvin
- D) It doesn't matter

3.112. In the ideal gas law, which variable represents the gas constant?

- A) n
- B) R
- C) V
- D) T

3.117. The SI unit for pressure

- A) bar
- B) kPa
- C) atm
- D) Pa

3.123. What Is The Formula For Boyle's Law?

- A)  $p_i \times V_i = p_f \times V_f$
- B)  $p \times V = \text{zero}$
- C)  $p/V = \text{const}$
- D)  $p/T = \text{const}$

3.124. In the Boyle's Law Formula  $V_i$  is the

- A) initial pressure
- B) initial volume
- C) final pressure
- D) final volume


3.125. In the Boyle's Law Formula  $p_f$  is the

- A) initial pressure
- B) initial volume
- C) final pressure
- D) final volume

3.126. True or false? Boyle's Law means: the volume of a mass of gas is inversely proportional to its pressure.

- A) True
- B) False

3.127. What is the correct option?



- A)  $T_1 = T_2$
- B)  $T_1 > T_2$
- C)  $T_1 < T_2$

3.128. If 22.5 dm<sup>3</sup> of nitrogen at 758 Hgmm are compressed to 737 Hgmm at constant temperature. What is the new volume in m<sup>3</sup>?

Solution:

$$p_1 \times V_1 = p_2 \times V_2$$

$$V_2 = (p_1 \times V_1) / p_2$$

$$V_1 = 22.5 / 1000 = 0.0225 \text{ m}^3$$

$$1 \text{ [mmHg]} = 133.3224 \text{ [Pa]}$$

$$p_1 = 758 \text{ Hgmm} = 101058,38 \text{ Pa}$$

$$p_2 = 737 \text{ Hgmm} = 98258,61 \text{ Pa}$$

$$V_2 = (101058,38 \times 0.0225) / 98258,61 = 0.023 \text{ m}^3$$

3.129. 2000 cm<sup>3</sup> of a gas at standard temperature and pressure is compressed to 1473 mL. What is the new pressure of the gas?

Solution:

$$p_1 = 1 \text{ atm}$$

$$V_1 = 2000 \text{ cm}^3 = 2 \text{ dm}^3 = 0.002 \text{ m}^3$$

$$V_2 = 1473 \text{ mL} = 1.473 \text{ dm}^3 = 0.001473 \text{ m}^3$$


$$p_2 = ?$$

$$p_1 \times V_1 = p_2 \times V_2$$

$$p_2 = [p_1 \times V_1] / V_2$$

$$p_2 = [1 \times 0.002] / 0.001473 = 1.358 \text{ atm}$$

3.132. What is isochoric process in the diagram?



A) (3) → (1)

B) (2) → (3)

C) (2) → (1)

D) (1) → (2)

3.134. The symbol of the gas constant

A) k

B) R

C) f

D) n

3.135. The units of the gas constant

- A) bar/K
- B) Pa/K
- C) J/mol·K
- D) cal/mol

3.136. The symbol of the Boltzmann constant

- A) B
- B) z
- C) c
- D) k

3.137. Which is the correct answer?

- A)  $k = 8.31 \text{ J} / \text{K}^{-1}$
- B)  $k = 1.3806 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$
- C)  $k = 6 \times 10^{23}$
- D)  $k = 1.3806 \times 10^{-26} \text{ J} \cdot \text{K}^{-1}$

3.138. R is related to the Boltzmann constant (k) by

- A)  $R = m/M \cdot k \cdot N_A$
- B)  $R = n \cdot k \cdot N_A$
- C)  $R = k \cdot N_A$
- D)  $R = N_A / k$

3.142. Fill up the missing spots in the table below!

Type of Gas Law	Formula	Description
1.	2. $p_1 \times V_1 = p_2 \times V_2$	3.
4.	5.	6.
7.	8.	9.
10. Charles' Law	11.	12.
13.	14. $pV = nRT$	

- A)  $V_1 / T_1 = V_2 / T_2$
- B) Combined Law
- C)  $p_1 / T_1 = p_2 / T_2$
- D) Obtained by combining Boyle's Law, Charles' Law and Gay-Lussac's Law
- E) Boyle's Law
- F) At constant P, as volume increases, temperature increases
- G)  $(p_1 \times V_1) / T_1 = (p_2 \times V_2) / T_2$
- H) At constant T, as pressure increases, volume decreases
- I) Gay - Lussac's Law
- J) At constant V, as pressure increases, temperature increases

3.148. In the gas laboratory is an air compressor. The tank of the leaky air compressor originally holds 100 dm<sup>3</sup> of air at 28 °C and 207 kPa. During a compression process, 3 grams of air is lost, the remaining air occupies 47 L at 490 kPa. What is the temperature of the remaining air?

*Solution:*

$$p_1 = 207 \text{ kPa} = 207000 \text{ Pa}$$

$$R = 287 \text{ J/ kg} \times \text{K}$$

$$T_1 = 28 \text{ }^\circ\text{C} = 28 + 273 = 301 \text{ K}$$

$$V_1 = 100 \text{ dm}^3 = 0.1 \text{ m}^3$$

$$p_2 = 490 \text{ kPa} = 490000 \text{ Pa}$$

$$V_2 = 47 \text{ L} = 0.047 \text{ m}^3$$

$$\rho_1 = p_1 / RT_1 = 207000 / (287 \times 301) = 2.396 \text{ kg/m}^3$$

$$m_1 = \rho_1 \times V_1 = 2.396 \times 0.1 = 0.2396 \text{ kg}$$

$$\rho_2 = p_2 / RT_2$$

$$\rho_2 = m_2 / V_2 = [0.2396 - 0.003] / 0.047 = 5.034 \text{ kg /m}^3$$

$$5.034 = 490000 / [287 \times T_2]$$

$$1444.758 T_2 = 490000$$

$$T_2 = 339.16 \text{ K}$$

3.153. Calculate the density, the specific weight and the volume of chloride gas at 27°C and pressure of 750 000 N/m<sup>2</sup>.

$$T = 27 + 273 = 300 \text{ K}$$

$$R = 118 \text{ J / mol K}$$

$$p = 750 \text{ 000 N/m}^2$$

*Solution:*

$$\rho = p / R \cdot T$$

$$\rho = 750000 / (118 \times 300) = 21.186 \text{ kg/m}^3$$

$$\gamma = \rho \times g = 21.186 \times 9.81 = 207.83 \text{ N/m}^3$$

$$V = 1 / \rho = 1 / 21.186 = 0.0472 \text{ m}^3 / \text{kg}$$

3.168. How to Convert Kelvin to Celsius?

A)  $T(^{\circ}\text{C}) = T(\text{K}) - 273.15$

B)  $T(^{\circ}\text{C}) = T(\text{K}) - 273$

C)  $T(^{\circ}\text{C}) = T(\text{K}) + 273.15$

D)  $T(^{\circ}\text{C}) = T(\text{K}) + 273$

3.169. What are extensive properties in thermodynamics?

- A) density, mass, length
- B) mass, volume, length
- C) boiling point, mass, density
- D) mass, volume, shape

3.170. The mass is 600 g and the volume is 200 cm<sup>3</sup>. Determine the specific volume!

- A) 3 g/cm<sup>3</sup>
- B) 0.33 g/cm<sup>3</sup>
- C) 3.33 x 10<sup>-4</sup> m<sup>3</sup>/kg
- D) 0.33 m<sup>3</sup>/kg

Solution:

$$m = 600 \text{ g} = 0.6 \text{ kg}$$

$$V = 200 \text{ cm}^3 = 0.0002 \text{ m}^3$$

$$\text{specific volume} = 0.0002 / 0.6 = 3.33 \times 10^{-4} \text{ m}^3/\text{kg}$$

4.7. The weight density is given by

- A)  $w = \gamma \times g$
- B)  $w = \rho \times g$
- C)  $\rho = w \times g$
- D)  $\gamma = w \times g$

4.20. Absolute pressure

- A)  $p_{\text{abs}} = p_{\text{atm}} - p_{\text{gage}}$
- B)  $p_{\text{abs}} = p_{\text{atm}} + p_{\text{gage}}$
- C)  $p_{\text{abs}} = p_{\text{atm}}$
- D)  $p_{\text{abs}} = p_{\text{gage}}$

4.21. Pressure head

- A)  $Z = p \times (\rho \times g)$
- B)  $Z = p / (\rho \times g)$
- C)  $Z = \gamma \times h$
- D)  $Z = \rho \times g \times h$

4.23. Which of the following is the hydrostatic equation?

- A)  $P = \rho g d$
- B)  $P = \rho g h$
- C)  $P / (\rho g) = d$
- D)  $\rho = P / (g d)$

4.54. A U-tube about half filled with liquid, with both ends of the tube open, the liquid is

- A) at the same height in each leg
- B) at the different height in each leg